



## VI. PREVALENCE OF USE BY GENDER

### **Prevalence Rates by Gender: Lifetime and Monthly Use**

• Grade 6 .....	94
• Grade 7 .....	95
• Grade 8 .....	96
• Grade 9 .....	97
• Grade 10 .....	98
• Grade 11 .....	99
• Grade 12 .....	100

**Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**6th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	8.3	6.3	7.3
Smokeless tobacco*	4.0	1.0	2.5
Cigars*	2.9	0.9	1.9
Pipe*	1.7	0.9	1.3
Alcohol*	19.9	14.8	17.4
Marijuana*	3.9	2.3	3.1
Cocaine*	0.7	0.4	0.6
Crack	0.6	0.5	0.6
Inhalants*	2.2	1.4	1.8
Amphetamines	0.5	0.3	0.4
Methamphetamines	0.6	0.4	0.5
Heroin	0.5	0.3	0.4
Steroids*	0.8	0.4	0.6
Ecstasy*	0.5	0.3	0.4
Tranquilizers	0.6	0.4	0.5
Hallucinogens	0.5	0.4	0.4
Prescription painkillers	1.2	1.0	1.1
Prescription drugs*	1.2	0.9	1.0
Over the counter drugs	1.2	1.0	1.1

NOTES: \* Differences between males and females are statistically significant (p < .05).

**Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**6th Grade**

DRUG	Male	Female	Indiana
Cigarettes	2.7	2.6	2.6
Smokeless tobacco*	1.5	0.4	1.0
Cigars*	0.9	0.4	0.7
Pipe*	0.8	0.5	0.6
Alcohol*	5.8	4.9	5.3
Binge drinking**	6.3	5.4	5.8
Marijuana*	1.9	1.4	1.6
Cocaine*	0.4	0.2	0.3
Crack*	0.3	0.2	0.3
Inhalants	0.9	0.8	0.9
Amphetamines	0.3	0.2	0.2
Methamphetamines	0.3	0.2	0.3
Heroin	0.3	0.1	0.2
Steroids*	0.3	0.1	0.2
Ecstasy*	0.3	0.2	0.2
Tranquilizers*	0.3	0.2	0.3
Hallucinogens	0.3	0.2	0.3
Prescription painkillers	0.6	0.6	0.6
Prescription drugs	0.6	0.6	0.6
Over the counter drugs	0.7	0.7	0.7

NOTES: \* Differences between males and females are statistically significant (p < .05).  
# Binge drinking during the past two weeks.

### Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 7th Grade

DRUG	Male	Female	Indiana
Cigarettes	13.7	12.7	13.2
Smokeless tobacco*	7.7	2.5	5.1
Cigars*	6.1	2.5	4.3
Pipe*	4.2	2.3	3.2
Alcohol	27.4	24.7	26.1
Marijuana*	8.3	6.3	7.3
Cocaine*	1.5	0.7	1.1
Crack	1.4	0.9	1.1
Inhalants	3.2	3.2	3.2
Amphetamines	1.3	0.8	1.0
Methamphetamines*	1.2	0.8	1.0
Heroin	1.0	0.6	0.8
Steroids*	1.4	0.6	1.0
Ecstasy*	1.3	0.8	1.1
Tranquilizers	1.1	0.8	0.9
Hallucinogens*	1.2	0.6	0.9
Prescription painkillers	2.8	2.5	2.7
Prescription drugs	2.4	2.3	2.4
Over the counter drugs*	2.6	2.9	2.8

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).

### Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 7th Grade

DRUG	Male	Female	Indiana
Cigarettes*	5.9	6.1	6.0
Smokeless tobacco*	3.4	1.0	2.2
Cigars*	2.1	1.0	1.6
Pipe*	1.7	1.2	1.5
Alcohol*	9.9	10.5	10.2
Binge drinking#	7.8	8.0	7.9
Marijuana*	4.5	3.7	4.1
Cocaine*	0.9	0.4	0.6
Crack	0.9	0.5	0.7
Inhalants	1.4	1.5	1.4
Amphetamines	0.7	0.4	0.5
Methamphetamines*	0.6	0.4	0.5
Heroin	0.6	0.3	0.4
Steroids*	0.7	0.3	0.5
Ecstasy*	0.6	0.4	0.5
Tranquilizers	0.5	0.4	0.5
Hallucinogens	0.6	0.4	0.5
Prescription painkillers*	1.5	1.4	1.5
Prescription drugs	1.4	1.5	1.4
Over the counter drugs*	1.4	1.8	1.6

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).  
# Binge drinking during the past two weeks.

**Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**8th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	20.3	20.9	20.6
Smokeless tobacco*	13.0	4.4	8.6
Cigars*	11.1	5.3	8.1
Pipe*	7.2	4.8	6.0
Alcohol*	36.5	38.8	37.7
Marijuana*	15.4	13.9	14.6
Cocaine*	2.3	1.8	2.0
Crack*	1.7	1.6	1.7
Inhalants*	5.2	5.7	5.4
Amphetamines	2.4	2.3	2.3
Methamphetamines	1.6	1.4	1.5
Heroin*	1.4	1.0	1.2
Steroids*	1.8	0.8	1.3
Ecstasy*	2.5	1.9	2.2
Tranquilizers	1.9	1.9	1.9
Hallucinogens*	2.6	2.1	2.3
Prescription painkillers*	5.3	5.9	5.6
Prescription drugs*	4.4	5.1	4.8
Over the counter drugs*	4.2	5.7	5.0

NOTES: \* Differences between males and females are statistically significant (p < .05).

**Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**8th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	9.4	10.6	10.0
Smokeless tobacco*	6.2	1.6	3.9
Cigars*	4.3	2.1	3.2
Pipe*	3.3	2.7	3.0
Alcohol*	14.9	18.4	16.7
Binge drinking**	10.5	11.7	11.1
Marijuana*	8.6	8.1	8.3
Cocaine	1.3	0.8	1.0
Crack*	1.0	0.7	0.9
Inhalants*	2.1	2.5	2.3
Amphetamines	1.3	1.1	1.2
Methamphetamines	1.0	0.8	0.9
Heroin	0.8	0.5	0.7
Steroids*	1.0	0.4	0.7
Ecstasy*	1.3	0.9	1.1
Tranquilizers	1.0	1.0	1.0
Hallucinogens*	1.4	1.0	1.2
Prescription painkillers*	2.8	3.3	3.0
Prescription drugs*	2.4	3.0	2.7
Over the counter drugs*	2.1	3.2	2.7

NOTES: \* Differences between males and females are statistically significant (p < .05).  
# Binge drinking during the past two weeks.

### Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 9th Grade

DRUG	Male	Female	Indiana
Cigarettes	27.3	27.1	27.2
Smokeless tobacco*	19.8	6.7	13.1
Cigars*	20.6	9.4	14.9
Pipe*	12.6	8.3	10.4
Alcohol*	46.2	49.2	47.7
Marijuana*	24.9	21.1	23.0
Cocaine*	3.7	2.2	2.9
Crack*	2.4	1.5	1.9
Inhalants	7.0	6.8	6.9
Amphetamines	4.4	4.1	4.3
Methamphetamines*	2.2	1.5	1.9
Heroin*	2.0	1.2	1.6
Steroids*	2.2	0.8	1.5
Ecstasy*	3.8	3.0	3.4
Tranquilizers	3.4	3.2	3.3
Hallucinogens*	4.3	2.9	3.6
Prescription painkillers	9.6	9.2	9.4
Prescription drugs*	8.1	8.3	8.2
Over the counter drugs*	6.5	7.9	7.2

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).  
# Binge drinking during the past two weeks.

### Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 9th Grade

DRUG	Male	Female	Indiana
Cigarettes	14.1	14.6	14.4
Smokeless tobacco*	10.3	2.1	6.2
Cigars*	8.5	3.5	6.0
Pipe*	5.6	3.9	4.8
Alcohol*	22.7	25.0	23.8
Binge drinking#	15.5	15.1	15.3
Marijuana*	14.1	11.8	13.0
Cocaine*	1.9	0.9	1.4
Crack*	1.3	0.7	1.0
Inhalants	2.2	2.3	2.2
Amphetamines	1.9	1.9	1.9
Methamphetamines*	1.0	0.7	0.9
Heroin*	1.0	0.6	0.8
Steroids*	1.1	0.4	0.8
Ecstasy*	1.7	1.2	1.4
Tranquilizers	1.6	1.6	1.6
Hallucinogens*	1.9	1.2	1.5
Prescription painkillers	4.5	4.9	4.7
Prescription drugs	4.0	4.6	4.3
Over the counter drugs*	3.0	4.1	3.5

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).  
# Binge drinking during the past two weeks.

**Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**10th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	34.3	31.4	32.8
Smokeless tobacco*	26.7	8.3	17.2
Cigars*	28.5	13.8	20.9
Pipe*	18.3	11.9	15.0
Alcohol*	54.7	56.3	55.5
Marijuana*	32.8	28.0	30.3
Cocaine*	5.0	2.7	3.8
Crack*	2.9	1.3	2.0
Inhalants*	8.0	7.0	7.4
Amphetamines*	6.4	5.5	5.9
Methamphetamines*	2.8	1.6	2.2
Heroin*	2.6	1.2	1.9
Steroids*	2.5	0.8	1.7
Ecstasy*	5.4	3.5	4.4
Tranquilizers	4.6	4.0	4.3
Hallucinogens*	6.5	3.7	5.1
Prescription painkillers*	13.3	11.9	12.6
Prescription drugs	11.5	10.9	11.2
Over the counter drugs	9.0	8.7	8.8

NOTES: \* Differences between males and females are statistically significant (p < .05).

**Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**10th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	18.5	16.1	17.2
Smokeless tobacco*	14.0	2.3	7.9
Cigars*	12.8	4.9	8.7
Pipe*	8.0	5.2	6.6
Alcohol	28.9	27.9	28.4
Binge drinking**	19.8	16.6	18.1
Marijuana*	18.6	14.4	16.4
Cocaine*	2.1	1.0	1.6
Crack*	1.4	0.6	1.0
Inhalants*	2.4	1.9	2.1
Amphetamines*	2.6	2.1	2.4
Methamphetamines*	1.3	0.6	0.9
Heroin*	1.3	0.5	0.9
Steroids*	1.3	0.4	0.8
Ecstasy*	2.2	1.3	1.7
Tranquilizers	2.1	1.8	2.0
Hallucinogens*	2.5	1.3	1.9
Prescription painkillers*	6.2	5.5	5.8
Prescription drugs	5.4	5.1	5.2
Over the counter drugs	3.6	3.6	3.6

NOTES: \* Differences between males and females are statistically significant (p < .05).  
# Binge drinking during the past two weeks.

### Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 11th Grade

DRUG	Male	Female	Indiana
Cigarettes*	40.0	36.3	38.1
Smokeless tobacco*	31.2	9.9	20.0
Cigars*	37.0	18.0	27.0
Pipe*	24.3	16.3	20.1
Alcohol*	60.7	60.5	60.6
Marijuana*	38.8	32.3	35.4
Cocaine*	7.1	3.3	5.1
Crack*	3.5	1.5	2.5
Inhalants*	9.0	6.9	7.9
Amphetamines*	8.5	7.2	7.8
Methamphetamines*	3.6	2.0	2.7
Heroin*	3.3	1.2	2.2
Steroids*	3.2	0.9	2.0
Ecstasy*	7.1	4.4	5.7
Tranquilizers	5.4	5.0	5.2
Hallucinogens*	8.5	4.7	6.5
Prescription painkillers*	16.8	13.2	14.9
Prescription drugs*	14.3	12.4	13.3
Over the counter drugs*	10.3	8.8	9.5

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).

### Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

#### 11th Grade

DRUG	Male	Female	Indiana
Cigarettes*	22.1	19.5	20.7
Smokeless tobacco*	15.6	2.7	8.8
Cigars*	16.7	6.0	11.1
Pipe*	10.0	6.8	8.3
Alcohol*	33.3	30.3	31.7
Binge drinking**	23.8	18.2	20.8
Marijuana*	21.6	15.8	18.6
Cocaine*	2.7	1.0	1.8
Crack*	1.8	0.6	1.2
Inhalants*	2.1	1.5	1.8
Amphetamines*	2.7	2.4	2.6
Methamphetamines*	1.9	0.7	1.3
Heroin*	1.7	0.6	1.1
Steroids*	1.6	0.4	1.0
Ecstasy*	2.4	1.2	1.8
Tranquilizers	2.2	1.7	2.0
Hallucinogens*	3.1	1.4	2.2
Prescription painkillers*	6.8	5.1	5.9
Prescription drugs*	6.0	5.3	5.6
Over the counter drugs*	3.5	3.2	3.3

NOTES: \* Differences between males and females are statistically significant ( $p < .05$ ).

# Binge drinking during the past two weeks.

**Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**12th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	45.1	39.2	42.1
Smokeless tobacco*	36.6	10.7	23.6
Cigars*	48.1	23.9	35.9
Pipe*	32.5	21.8	27.1
Alcohol	65.6	65.1	65.4
Marijuana*	42.2	35.1	38.6
Cocaine*	8.1	3.8	5.9
Crack*	3.7	1.5	2.6
Inhalants*	9.5	6.1	7.8
Amphetamines*	9.8	6.9	8.3
Methamphetamines*	3.7	1.9	2.8
Heroin*	3.4	1.4	2.4
Steroids*	3.4	0.9	2.1
Ecstasy*	8.4	4.9	6.7
Tranquilizers*	6.1	4.8	5.4
Hallucinogens*	9.8	4.6	7.2
Prescription painkillers*	19.0	13.1	16.0
Prescription drugs*	16.5	12.7	14.6
Over the counter drugs*	11.7	7.8	9.7

NOTES: \* Differences between males and females are statistically significant (p < .05).

**Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011**  
(Values expressed as percentages)

**12th Grade**

DRUG	Male	Female	Indiana
Cigarettes*	27.9	21.4	24.6
Smokeless tobacco*	19.2	2.4	10.7
Cigars*	25.1	8.4	16.7
Pipe*	15.1	9.9	12.5
Alcohol*	40.3	35.4	37.8
Binge drinking**	29.9	23.0	26.4
Marijuana*	23.7	16.0	19.8
Cocaine*	2.9	1.2	2.0
Crack*	1.7	0.6	1.2
Inhalants*	2.4	1.0	1.7
Amphetamines*	3.4	2.1	2.7
Methamphetamines*	1.8	0.8	1.3
Heroin*	1.8	0.6	1.2
Steroids*	1.7	0.4	1.1
Ecstasy*	3.0	1.2	2.1
Tranquilizers	2.4	1.6	2.0
Hallucinogens*	3.6	1.3	2.4
Prescription painkillers*	8.3	4.9	6.6
Prescription drugs*	6.9	4.9	5.9
Over the counter drugs*	4.0	2.4	3.2

NOTES: \* Differences between males and females are statistically significant (p < .05).  
# Binge drinking during the past two weeks.